

Anti-CP47 subunit of Photosystem II antibody

Catalog: PHY0058A

Product Information

Description:	Rabbit polyclonal antibody
Background:	CP47 is one of the components of the core complex of photosystem II (PSII). It binds chlorophyll and helps catalyze the primary light-induced photochemical processes of PSII. CP47 is encoded by ATCG00680.
Synonyms:	PsbB
Immunogen:	KLH-conjugated synthetic peptide of CP47 subunit derived from <i>Arabidopsis thaliana</i> ATCG00680.
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected/apparent MW:	56 / 47 kDa
Confirmed Reactivity:	<i>Arabidopsis thaliana</i> , <i>Glycine max</i> , <i>Zea mays</i> , <i>Oryza sativa Indica Group</i> , <i>Hordeum vulgare subsp. Vulgare</i>
Predicted Reactivity:	Among 25 analyzed species, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in <i>Physcomitrella patens</i> , <i>Spinacia oleracea</i> , <i>Medicago truncatula</i> ,

Research Use Only

Gossypium raimondii, *Spinacia oleracea*, *Brassica napus*, *Nicotiana tabacum*, *Setaria viridis*, *Populus trichocarpa*, *Vitis vinifera*, *Cucumis sativus*, *Sorghum bicolor*, *Nicotiana tabacum*, and 80-99% homologues with the sequence in *Triticum aestivum*, *Chlamydomonas reinhardtii*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



1-3 is thylakoid membrane protein from *Arabidopsis thaliana* leaf containing 0.1 µg, 0.25 µg, and 0.5 µg of chlorophyll, respectively.

Electrophoresis: 15% SDS-Urea-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1h.

Blocking: 5% skim milk at RT or 4°C for 1h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:20000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.