

# Anti-CP47 subunit of Photosystem II antibody

Catalog: PHY0058A

## Product Information

<b>Description:</b>	Rabbit polyclonal antibody
<b>Background:</b>	CP47 is one of the components of the core complex of photosystem II (PSII). It binds chlorophyll and helps catalyze the primary light-induced photochemical processes of PSII. CP47 is encoded by ATCG00680.
<b>Synonyms:</b>	PsbB
<b>Immunogen:</b>	KLH-conjugated synthetic peptide of CP47 subunit derived from <i>Arabidopsis thaliana</i> ATCG00680.
<b>Form:</b>	Lyophilized
<b>Quantity:</b>	150 µg
<b>Purification:</b>	Immunogen affinity purified
<b>Reconstitution:</b>	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
<b>Stability &amp; Storage:</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
<b>Shipping:</b>	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

## Application Information


<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
<b>Expected/apparent MW:</b>	56 / 47 kDa
<b>Confirmed Reactivity:</b>	<i>Arabidopsis thaliana</i> , <i>Glycine max</i> , <i>Zea mays</i> , <i>Oryza sativa Indica Group</i> , <i>Hordeum vulgare subsp. Vulgare</i>
<b>Predicted Reactivity:</b>	Among 25 analyzed species, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in <i>Physcomitrella patens</i> , <i>Spinacia oleracea</i> , <i>Medicago truncatula</i> ,

Research Use Only

*Gossypium raimondii*, *Spinacia oleracea*, *Brassica napus*, *Nicotiana tabacum*, *Setaria viridis*, *Populus trichocarpa*, *Vitis vinifera*, *Cucumis sativus*, *Sorghum bicolor*, *Nicotiana tabacum*, and 80-99% homologues with the sequence in *Triticum aestivum*, *Chlamydomonas reinhardtii*.

For more species homologues information, please contact tech support at [tech@phytoab.com](mailto:tech@phytoab.com).

## Application Example



1-3 is thylakoid membrane protein from *Arabidopsis thaliana* leaf containing 0.1 µg, 0.25 µg, and 0.5 µg of chlorophyll, respectively.

**Electrophoresis:** 15% SDS-Urea-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1h.

**Blocking:** 5% skim milk at RT or 4°C for 1h.

**Primary antibody:** 1:2000 dilution overnight at 4°C.

**Secondary antibody:** 1:20000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.