

# Anti-PsbR subunit of Photosystem II antibody

Catalog: PHY0110A

#### **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** PsbR is the subunit of photosystem II (PSII) with a molecular mass of 10 kDa.

This subunit appears to be involved in the stable assembly of PSII, particularly

that of the oxygen-evolving complex subunit PsbP. Mutants defective in

accumulation of PsbR have reduced amounts of subunits PsbP and PsbQ in

PSII.

Synonyms: PsbR, Photosystem II 10 kDa polypeptide, chloroplastic, photosystem II subunit

R.

**Immunogen:** KLH-conjugated synthetic peptide of PsbR derived from *Arabidopsis thaliana* 

AT1G79040.

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70℃ as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected/apparent MW:** 14 / 10 kDa

Confirmed Reactivity: Arabidopsis thaliana




#### **Predicted Reactivity:**

Among 25 analyzed species, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Brassica napus, Brassica rapa, Gossypium raimondii,*Chlamydomonas reinhardtii, and 80-99% homologues with the sequence in Cucumis sativus, Nicotiana tabacum, Nicotiana tabacum, Solanum lycopersicum, Solanum tuberosum, Vitis vinifera, Populus trichocarpa.

For more species homologues information, please contact tech support at tech@phytoab.com.

### **Application Example**



1-3 is thylakoid membrane protein from WT of Arabidopsis thaliana leaf containing 1  $\mu$ g, 2  $\mu$ g, and 5  $\mu$ g of chlorophyll, respectively.

4-6 is thylakoid membrane protein from PSII mutant of *Arabidopsis thaliana* leaf containing 1 μg, 2 μg, and 5 μg of chlorophyll, respectively.

Electrophoresis: Tricine-SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

**Primary antibody:** 1:1000 dilution overnight at  $4^{\circ}$ C.

Secondary antibody: 1:20000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.