

## Anti-Ribosomal protein S4, mitochondrial antibody

Catalog: PHY2145A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** RPS4 is one of the primary rRNA binding proteins, it binds directly to 16S

rRNA where it nucleates assembly of the body of the 30S subunit.

Synonyms: RPS4, MITOCHONDRIAL RIBOSOMAL PROTEIN S4, RIBOSOMAL

PROTEIN US4M, US4M

**Immunogen:** KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from

Arabidopsis thaliana RPS4 (ATMG00290 / AT2G07734).

Form: Lyophilized

**Quantity**: 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of sterile 1XPBS (PH=7.4).

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4 °C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 43 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide

used for immunization is 100% homologues with the sequence

in Brassica rapa, Brassica napus, and 80-99% homologues with the


sequence in Populus trichocarpa, Gossypium raimondii, Vitis

vinifera.



For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 47 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.