

## **Anti-Vacuolar-sorting protein BRO1 antibody**

Catalog: PHY7736S

## **Product Information**

**Description:** Rabbit polyclonal antibody

Background: ALIX mediates endosomal protein trafficking and turnover through the

ESCRT/MVB pathway. One target is the phosphate transporter PHT1;1. Modulates ABA-mediated inhibition of plant growth and stomatal aperture

(PMID:31363038).

Synonyms: ALIX, ALG-2 INTERACTING PROTEIN-X

**Immunogen:** KLH-conjugated synthetic peptide (16 aa from N terminal section) derived from

Arabidopsis thaliana ALIX (AT1G15130).

Form: Lyophilized

**Quantity:** 150 μg

Purification:

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

Reconstitution: Reconstitution with 150 µl of sterile water.

Serum

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to  $-70^{\circ}$ C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4 °C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 95 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used for


immunization is 80-99% homologues with the sequence in Brassica

rapa, Brassica napus, Cucumis sativus, Populus trichocarpa.



For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 60 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.