

Anti-Beta-amylase 5, C-terminal antibody

Catalog: PHY2278A

Product Information

Description:	Rabbit polyclonal antibody
Background:	The Arabidopsis (Arabidopsis thaliana) genome contains nine β -amylase
	(BAM) genes, some of which play important roles in starch hydrolysis.
	RAM1(BAM5) has beta-amylase activity. It is major cytosolic beta-amylase
	isoform in rosette leaves and inflorescences stems.
Synonyms:	RAM1, ARABIDOPSIS THALIANA BETA-AMYLASE, AT-BETA-AMY,
	ATBETA-AMY, BAM5, BETA-AMYLASE 5, BMY1, REDUCED BETA AMYLASE
	1
Immunogen:	KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from
	Arabidopsis thaliana RAM1 (AT4G15210).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 μl of sterile 1XPBS (PH=7.4).
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ C$ as supplied.
	6 months, -20 to -70 $^\circ$ C under sterile conditions after reconstitution.
	1 month, 2 to 8 $^\circ \!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!\!$
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	56 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used


Nesearc



for immunization is 100% homologues with the sequence in *Brassica napus*, *Brassica rapa*.

For more species homologues information, please contact tech support at <u>tech@phytoab.com</u>.

Application Example



PHY2278A

Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa. **Electrophoresis:** 12% SDS-PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4° for 1 h.

Primary antibody: 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.