

Anti-Beta-amylase 5 antibody

Catalog: PHY2277A

Product Information

Description: Rabbit polyclonal antibody

Background: The Arabidopsis (*Arabidopsis thaliana*) genome contains nine β-amylase

(BAM) genes, some of which play important roles in starch hydrolysis.

RAM1(BAM5) has beta-amylase activity. It is major cytosolic beta-amylase

isoform in rosette leaves and inflorescences stems.

Synonyms: RAM1, ARABIDOPSIS THALIANA BETA-AMYLASE, AT-BETA-AMY,

ATBETA-AMY, BAM5, BETA-AMYLASE 5, BMY1, REDUCED BETA AMYLASE

1

Immunogen: KLH-conjugated synthetic peptide (16 aa from N terminal section) derived from

Arabidopsis thaliana RAM1 (AT4G15210).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 μl of sterile 1XPBS (PH=7.4).

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4 °C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 56 kDa


Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used



for immunization is 80-99% homologues with the sequence in *Brassica* napus, *Brassica* rapa, *Hordeum* vulgare.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.