

## Anti-Oligouridylate-binding protein 1C, C-terminal antibody

Catalog: PHY2111A

## **Product Information**

Description:	Rabbit polyclonal antibody
Background:	Low oxygen stress dynamically regulates the translation of cellular mRNAs as a
	means of energy conservation in seedlings of <i>Arabidopsis thaliana</i> . UBP1C is
	heterogeneous nuclear ribonucleoprotein (hnRNP)-like protein that acts as a
	component of the pre-mRNA processing machinery.
Synonyms:	UBP1C, F OLIGOURIDYLATE BINDING PROTEIN 1C
Immunogen:	KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from
	Arabidopsis thaliana UBP1C (AT3G14100).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
<b>Reconstitution:</b>	Reconstitution with 150 $\mu$ l of sterile 1 $ imes$ PBS (PH=7.4).
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb C$ as supplied.
	6 months, -20 to -70 $^\circ$ C under sterile conditions after reconstitution.
	1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\mathrm{C}$ . Upon receipt, store it immediately at the
	temperature recommended above.


## **Application Information**

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	47 kDa
Predicted Reactivity:	For more species homologues information, please contact tech
	support at <u>tech@phytoab.com</u> .

Research Use Only



## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$  for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only