

Anti-Nuclear cap-binding protein subunit 2 antibody

Catalog: PHY0198S

Product Information

Description: Rabbit polyclonal antibody

Background: CBP20 is a nuclear cap-binding protein that forms a heterodimeric complex

with ABH1 (AT2G13540) and is likely to participate in RNA metabolism. It recognizes and binds capped RNAs (m7GpppG-capped RNA) but requires

ABH1/CBP80 to stabilize the movement of its N-terminal loop and lock the CBC

into a high affinity cap-binding state with the cap structure.

Synonyms: CBP20, ATCBP20, CAP-BINDING PROTEIN 20

Immunogen: KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Arabidopsis thaliana CBP20 (AT5G44200).

Form: Lyophilized

Quantity: 150 μg

Purification: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70° C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8℃ under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected/apparent MW: 30 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in



Brassica napus, Brassica rapa.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 44 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.