

Anti-Mitochondrial outer membrane protein porin 3 antibody

Catalog: PHY0697S

Product Information

Description: Rabbit polyclonal antibody

Background: VDAC3 is a member of voltage-dependent anion channel (VDAC:

AT3G01280/VDAC1, AT5G67500/VDAC2, AT5G15090/VDAC3,

AT5G57490/VDAC4, AT5G15090/VDAC5). VDACs are reported to be

porin-type, beta-barrel diffusion pores. They are prominently localized in the outer mitochondrial membrane and are involved in metabolite exchange

between the organelle and the cytosol.

Synonyms: VDAC3, ARABIDOPSIS THALIANA VOLTAGE DEPENDENT ANION

CHANNEL 3, ATVDAC-3, ATVDAC3, VOLTAGE DEPENDENT ANION

CHANNEL 3

Immunogen: KLH-conjugated synthetic peptide of VDAC3 derived from Arabidopsis thaliana

AT5G15090.

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the



end user.

Expected/apparent MW: 29 kDa

Confirmed Reactivity: Coming soon

Predicted Reactivity: Among 25 analyzed species, the sequence of the synthetic peptide

used for immunization is 80-99% homologues with the sequence in

Brassica napus.

For more species homologues information, please contact tech

support at tech@phytoab.com.