

## Anti-Endoplasmin homolog, N-terminal antibody

Catalog: PHY7109S

## **Product Information**

Description:	Rabbit polyclonal antibody
Background:	SHD is an ortholog of GRP94, an ER-resident HSP90-like protein and is
	involved in regulation of meristem size and organization.
Synonyms:	SHD, ATHSP90-7, ATHSP90.7, HEAT SHOCK PROTEIN 90-7, HEAT SHOCK
	PROTEIN 90.7, HSP90, HSP90.7, SHEPHERD
Immunogen:	KLH-conjugated synthetic peptide (15 aa from N terminal section) derived from
	Arabidopsis thaliana SHD (AT4G24190).
Form:	Lyophilized
Quantity:	150 μg
Purification:	Serum
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .
<b>Reconstitution:</b>	Reconstitution with 150µl of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ\!\!\mathbb{C}$ as supplied.
	6 months, -20 to -70 $^\circ C$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.
Shipping:	The product is shipped at $4^\circ\! \mathbb{C}$ . Upon receipt, store it immediately at the
	temperature recommended above.
Application l	oformation

## **Application Information**


<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	94 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 8 <mark>0-99% homologu</mark> es with the sequence in
	Brassica napus, Brassica rapa, Spinacia oleracea, Vitis vinifera.
	For more species homologues information, please contact tech

**Research Use Only** 



support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$  for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

**Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.



**Research Use Only**