

Anti-Protein TOC75-3/4, chloroplastic, C-terminal antibody

Catalog: PHY0130A

Product Information

Description:	Rabbit polyclonal antibody
Background:	TOC75-III/IV are Component of the translocon outer membrane (TOC)
	complex.
Synonyms:	TOC75-III/IV, TRANSLOCON AT THE OUTER ENVELOPE MEMBRANE OF
	CHLOROPLASTS 75-III/IV
Immunogen:	KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from
	Arabidopsis thaliana TOC75-III (AT3G46740) and TOC75-IV (AT4G09080).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen Affinity Purified
Reconstitution:	Reconstitution with 150 μ l of sterile 1 $ imes$ PBS (PH=7.4).
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70 $^\circ \!\! \mathbb{C}$ as supplied.
	6 months, -20 to -70 $^\circ\!\!\!\!\!^\circ$ under sterile conditions after reconstitution.
	1 month, 2 to 8 $^{\circ}$ C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4 $^\circ\!\!\!\mathrm{C}$. Upon receipt, store it immediately at the
	temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	89 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Brassica</i>
	napus, Brassica rapa, Gossypium raimondii, Spinacia oleracea,
	Nicotiana tabacum, Solanum lycopersicum, Cucumis sativus, Vitis



vinifera, Medicago truncatula, Populus trichocarpa, Solanum tuberosum, Oryza sativa, Panicum virgatum, Zea mays, Setaria viridis, Sorghum bicolor, Hordeum vulgare, Triticum aestivum. The sequence of the synthetic peptide used for immunization is 93% (14 / 15) homologues with the sequence in TOC75-I (AT1G35860). For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 48 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only