

# Anti-Protein FAR-RED ELONGATED HYPOCOTYL 1 antibody

Catalog: PHY7914S

## **Product Information**

Description:	Rabbit polyclonal antibody
Background:	FHY1 is a positive regulator of photomorphogenesis in far-red light. Most FHY1
	abundant in young seedlings in the dark. It is localized in the nucleus and the
	cytoplasm. FHY1 is degraded through the 26S proteasome and regulated by
	PHYA.
Synonyms:	FHY1, FAR-RED ELONGATED HYPOCOTYL 1, FRY1, PAT3,
	PHYTOCHROME A SIGNAL TRANSDUCTION 3
Immunogen:	KLH-conjugated synthetic peptide (16 aa from N terminal section) derived from
	Arabidopsis thaliana FHY1 (AT2G37678).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .
Reconstitution:	Reconstitution with 150 µl of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70°C as supplied.
	6 months, -20 to -70°C under sterile conditions after reconstitution.
	1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the
	temperature recommended above.

# **Application Information**


<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	23 kDa



#### Predicted Reactivity:

Among species analyzed, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica napus*, *Brassica rapa*. For more species homologues information, please contact tech support at tech@phytoab.com.

### Application Example



PHY7914S

Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured with CCD camera.

**Research Use Only**