

Anti-VOLTAGE DEPENDENT ANION CHANNEL 2, C-terminal antibody

Catalog: PHY7980S

Product Information

Description: Rabbit polyclonal antibody

Background: VDAC2 is a member of voltage-dependent anion channel (VDAC:

AT3G01280/VDAC1, AT5G67500/VDAC2, AT5G15090/VDAC3,

AT5G57490/VDAC4, AT5G15090/VDAC5). VDACs are reported to be

porin-type, beta-barrel diffusion pores. They are prominently localized in the outer mitochondrial membrane and are involved in metabolite exchange

between the organelle and the cytosol.

Synonyms: VDAC2, ARABIDOPSIS THALIANA VOLTAGE DEPENDENT ANION

CHANNEL 2, ATVDAC2, VOLTAGE DEPENDENT ANION CHANNEL 2

Immunogen: KLH-conjugated synthetic peptide (17 aa from C terminal section) derived from

Arabidopsis thaliana VDAC2 (AT5G67500).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:5000)

Note: Optimal dilutions/concentrations should be determined by the

end user.



Expected / apparent MW: 33 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 80-99% homologues with the sequence in

Brassica napus, Brassica rapa, Gossypium raimondii.

For homologues with other species especially algae, please contact

tech support at tech@phytoab.com.