

## Anti-Thioredoxin-like protein CDSP32 antibody

Catalog: PHY2613S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** CDSP32 is a thioredoxin-like protein in plants and it is localized to chloroplasts

playing dual roles in stress adaptation and redox regulation.

Synonyms: CDSP32, ARABIDOPSIS THALIANACHLOROPLASTIC DROUGHT-

INDUCED STRESS PROTEIN OF32 KD, ATCDSP32, CHLOROPLASTIC

DROUGHT-INDUCED STRESS PROTEIN OF 32 K

**Immunogen:** KLH-conjugated synthetic peptide (14 aa from central section) derived from

Arabidopsis thaliana FLS1 (AT1G76080).

Form: Lyophilized

Quantity:  $150 \mu g$ Purification: Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 27 kDa

**Predicted Reactivity:** Among species analyzed, the sequence of the synthetic peptide

used for immunization is 85-99% homologues with the sequence in


Brassica napus, and Brassica rapa.

Research Use Only



For more species homologues information, please contact tech support at <a href="mailto:tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were

captured with CCD camera.