

## **Anti-SNAP33 antibody**

Catalog: PHY2857A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** SNAP33 is probably involved in cytokinesis and cell plate formation.

Synonyms: SNAP33, ATSNAP33, ATSNAP33B, SNP33, SOLUBLE

N-ETHYLMALEIMIDE-SENSITIVE FACTOR ADAPTOR PROTEIN 33

Immunogen: KLH-conjugated synthetic peptide (16 aa from central section) derived from

Arabidopsis thaliana SNAP33 (AT5G61210).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of sterile 1 x PBS (PH=7.4) containing 30% glycerol.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 34 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 100% homologues with the sequence in Glycine

max, Brassica napus, Populus trichocarpa, Brassica rapa,

Gossypium raimondii, 80-99% homologues with the sequence in


Spinacia oleracea, Nicotiana tabacum, Medicago truncatula, Oryza



sativa, Cucumis sativus, Solanum tuberosum, Solanum lycopersicum, Vitis vinifera, Triticum aestivum, Hordeum vulgare subsp. vulgare, Zea mays, Physcomitrium patens.

For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.