

Anti-Os01g0133400 protein antibody

Catalog: PHY4425A

Product Information

| Description: | Rabbit polyclonal antibody | |
|------------------------|--|--|
| Background: | Os12g02677500 | |
| Synonyms: | pGlcT1, PGLCT, OspGlcT1, OsPGLCT | |
| Immunogen: | KLH-conjugated synthetic peptide (15 aa from Central section) derived from | |
| | <i>Oryza sativa</i> pGlcT1 (Os01g0133400). | |
| Form: | Lyophilized | |
| Quantity: | 150 µg | |
| Purification: | Immunogen affinity purified | |
| Reconstitution: | Reconstitution with 150 μ l of sterile 1 $	imes$ PBS (PH=7.4). | |
| | "Note: please spin tube briefly prior to opening it to avoid any losses that might | |
| | occur from lyophilized material adhering to the cap or sides of the tube". | |
| Stability & | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. | |
| Storage: | 12 months from date of receipt, -20 to -70°C as supplied. | |
| | 6 months, -20 to -70°C under sterile conditions after reconstitution. | |
| | 1 month, 2 to 8°C under sterile conditions after reconstitution. | |
| Shipping: | The product is shipped at 4° C. Upon receipt, store it immediately at the | |
| | temperature recommended above. | |


Application Information

| Recommended Dilution: | Western Blot (1:1000-1:2000) |
|-------------------------|--|
| | Note: Optimal dilutions/concentrations should be determined by the |
| | end user. |
| Expected / apparent MW: | 57 kDa |
| Predicted Reactivity: | Among species analyzed, the sequence of the synthetic peptide used |
| | for immunization is 80-99% homologues with the sequence in |
| | Triticum aestivum, Hordeum vulgare, Medicago truncatula. |
| | For more species homologues information, please contact tech |
| | support at <u>tech@phytoab.com</u> . |

Research Use Only



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 48 kDa. **Electrophoresis:** 12% SDS-PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:1000 dilution overnight at 4°C. **Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only