

Anti-KAH8493045/Dof zinc finger protein antibody

Catalog: PHY6366A

Product Information

| Description: | Rabbit polyclonal antibody | |
|------------------------|--|--|
| Background: | KAH8493045/Dof zinc finger protein is a putative transcription factor that binds | |
| | specifically to a 5'-AA[AG]G-3' consensus core sequence. | |
| Synonyms: | KAH8493045, Dof zinc finger protein | |
| Immunogen: | KLH-conjugated synthetic peptide (15 aa from C terminal section) derived from | |
| | Populus deltoides KAH8493045. | |
| Form: | Lyophilized | |
| Quantity: | 150 µg | |
| Purification: | Immunogen affinity purified | |
| Reconstitution: | Reconstitution with 150 μ I of sterile 1XPBS (PH=7.4) containing 30% glycerol. | |
| | "Note: please spin tube briefly prior to opening it to avoid any losses that might | |
| | occur from lyophilized m <mark>aterial</mark> adhering to the cap or sides of the tube". | |
| Stability & | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. | |
| Storage: | 12 months from date of receipt, -20 to -70°C as supplied. | |
| | 6 months, -20 to -70°C under sterile conditions after reconstitution. | |
| | 1 month, 2 to 8°C under sterile conditions after reconstitution. | |
| Shipping: | The product is shipped at 4°C. Upon receipt, store it immediately at the | |
| | temperature recommended above. | |
| | | |


Application Information

| Recommended Dilution: | Western Blot (1:1000-1:2000) |
|------------------------------|--|
| | Note: Optimal dilutions/concentrations should be determined by the |
| | end user. |
| Expected / apparent MW: | 34 kDa |
| Predicted Reactivity: | Among species analyzed, the sequence of the synthetic peptide used |
| | for immunization is 88% homologues with the sequence in <i>Populus</i> |
| | trichocarpa. |
| | For more species homologues information, please contact tech |
| | support at tech@phytoab.com. |

Research Use Only



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only