

## **Anti-Enhanced Disease Susceptibility 1 antibody**

Catalog: PHY7124S

## **Product Information**

**Description:** Rabbit polyclonal antibody

Background: EDS1 (Enhanced Disease Susceptibility 1) is a crucial nucleocytoplasmic plant

protein acting as a central regulator and integrator of innate immunity. Primarily, it functions as a guardian and signal transducer for intracellular immune receptors (NLRs) that detect pathogen effector proteins. Upon effector recognition, EDS1 forms complexes (notably with PAD4) to activate robust

defense responses.

Synonyms: EDS1, ATEDS1, ENHANCED DISEASE SUSCEPTIBILITY 1

**Immunogen:** KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Arabidopsis thaliana EDS1 (AT3G48090).

Form: Lyophilized

Quantity:  $150 \mu g$ Purification: Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.


Expected / apparent MW: 72 kDa



**Predicted Reactivity:** 

For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 44 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.