

Anti-Probable chlorophyll(ide) b reductase NYC1, chloroplastic, C-terminal antibody

Catalog: PHY3041S

Product Information

Description: Rabbit polyclonal antibody

Background: NYC1 is a chlorophyll b reductase involved in the degradation of chlorophyll b

and LHCII (light harvesting complex II).

Synonyms: NYC1, NON-YELLOW COLORING 1

Immunogen: KLH-conjugated synthetic peptide (19 aa from C terminal section) derived from

Arabidopsis thaliana NYC1 (AT4G13250).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70 °C as supplied.

6 months, -20 to -70 °C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4 °C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 55 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used for

immunization is 100% homologues with the sequence in *Brassica rapa*, *Brassica napus*, *Populus trichocarpa*, *Nicotiana tabacum*, and 80-99%



homologues with the sequence in *Glycine max*, *Medicago truncatula*,


Cucumis sativus, Gossypium raimondii, Vitis vinifera, Solanum

lycopersicum, Solanum tuberosum, Spinacia oleracea, Zea mays,

Sorghum bicolor, Oryza sativa, Setaria viridis, Hordeum vulgare, Triticum aestivum.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4° C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.