

Anti-BRASSINAZOLE-RESISTANT 1 antibody

Catalog: PHY4609S

Product Information

Description:	Rabbit polyclonal antibody	
Background:	BRASSINAZOLE-RESISTANT 1 mediates downstream brassinosteroid-	
	regulated growth response and feedback inhibition of brassinosteroid (BR)	
	biosynthetic genes.	
Synonyms:	Os07g0580500, OsBZR1, Protein BZR1 homolog 1	
Immunogen:	KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from	
	<i>Oryza sativa</i> BZR1 (Os07g0580500).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at info@phytoab.com.	
Reconstitution:	Reconstitution with 150 µl of sterile water.	
	"Note: please spin tube briefly prior to op <mark>ening it to avoid any losses that might</mark>	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70°C as supplied.	
	6 months, -20 to -70°C under sterile conditions after reconstitution.	
	1 month, 2 to 8°C under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the	
	temperature recommended above.	

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	32 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 87-99% homologues with the sequence in <i>Zea</i>
	mays, Hordeum vulgare, Triticum aestivum, Setaria viridis, Panicum
	virgatum, Sorghum bicolor.

Research Use Only



For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



PHY4609S

Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 42 kDa. Electrophoresis: 12% SDS-PAGE Transfer: blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. Primary antibody: 1:1000 dilution overnight at 4 °C. Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000). Detection: using chemiluminescence substrate and image were captured with CCD camera.