

## **Anti-LIKE HETEROCHROMATIN PROTEIN 1 antibody**

Catalog: PHY7866S

## **Product Information**

**Description:** Rabbit polyclonal antibody

Background: LHP1 protein localized preferentially to euchromatic regions and not to

heterochromatic chromocenters. It is involved in euchromatin organization and

is required for epigenetic maintenance of the vernalized state.

Synonyms: TFL2 (TERMINAL FLOWER 2) (Primary Symbol), AtLHP1, LHP1 (LIKE

HETEROCHROMATIN PROTEIN 1)

**Immunogen:** KLH-conjugated synthetic peptide (16 aa from central section) derived from

Arabidopsis thaliana LHP1 (AT5G17690).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## Application Information

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.


Expected / apparent MW: 49 kDa

**Predicted Reactivity:** For more species homologues information, please contact tech

support at tech@phytoab.com.



## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the

peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4 °C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.