

## Anti-WRKY DNA-BINDING PROTEIN 33 antibody

Catalog: PHY1840A

## **Product Information**

Description:	Rabbit polyclonal antibody
Background:	WRKY33 is a key pathogen-responsive transcription factor in Arabidopsis. It
	activates phytoalexin biosynthesis and defense genes during fungal/bacterial
	infections. Phosphorylated by MPK3/MPK6 upon PAMP recognition, WRKY33
	confers resistance by regulating salicylic acid/jasmonate signaling.
Synonyms:	WRKY33, WRKY DNA-BINDING PROTEIN 33
Immunogen:	KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from
	Arabidopsis thaliana WRKY33 (AT2G38470).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
<b>Reconstitution:</b>	Reconstitution with 150 µl of st <mark>erile 1×PB</mark> S (PH=7.4) containing 30% glycerol.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70°C as supplied.
	6 months, -20 to -70°C under sterile conditions after reconstitution.
	1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the
	temperature recommended above.


## **Application Information**

<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	57 kDa
Predicted Reactivity:	For more species homologues information, please contact tech
	support at tech@phytoab.com.

Research Use Only



## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 51 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.



Research Use Onl