

Anti-LHCA5 antibody

Product Information

Description:	Rabbit polyclonal antibody
Background:	Lhca5 is a low abundant chlorophyll a/b-binding protein that has been found to
	associate with photoystem I (PSI) in sub-stoechiometric amounts. It is essential
	for the interaction between NDH and one copy of PSI in higher plants.
Synonyms:	PHOTOSYSTEM I LIGHT HARVESTING COMPLEX GENE 5, LHCA5
Immunogen:	KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from
	Arabidopsis thaliana Lhca5 (AT1G45474).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 μ l of sterile 1XPBS (PH=7.4) containing 30% glycerol.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70°C as supplied.
	6 months, -20 to -70°C under sterile conditions after reconstitution.
	1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the
	temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	28 / 23 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used
	for immunization is 100% homologues with the sequence in <i>Glycine</i>
	max, Populus trichocarpa, Hordeum vulgare, Triticum aestivum, and
	80-99% homologues with the sequence in <i>Medicago truncatula,</i>



Brassica napus, Brassica rapa, Gossypium raimondii, Spinacia oleracea, Nicotiana tabacum, Vitis vinifera, Setaria viridis, Panicum virgatum, Oryza sativa, Solanum tuberosum, Solanum lycopersicum, Zea mays, Sorghum bicolor.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only