

Anti-Suppressor of TIC40 2 antibody

Catalog: PHY3992S

Product Information

Description: Rabbit polyclonal antibody

Background: STIC2 was identified in a screen for suppressors of chloroplast protein import

defect in tic40. It participates with ALB4 in thylakoid protein targeting and may

function with specific subset of thylakoidal proteins.

Synonyms: STIC2, Suppressor of TIC40 2

Immunogen: KLH-conjugated synthetic peptide (17 aa from C terminal section) derived from

Arabidopsis thaliana STIC2 (AT2G24020).

Form: Lyophilized

Quantity: $150 \mu g$ Purification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected/apparent MW: 20 kDa

Predicted Reactivity: Among 25 analyzed species, the sequence of the synthetic peptide

used for immunization is 94% homologues with the sequence in

Brassica napus, Brassica rapa, Medicago truncatula.


The sequence of the synthetic peptide used for immunization is 94%



homologues with the sequence in AT4G30620 (STCL).

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 42 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.