

# Anti-S-ADENOSYLMETHIONINE DECARBOXYLASE

## antibody

Catalog: PHY7905S

### **Product Information**

Description:	Rabbit polyclonal antibody
Background:	SAMDC is a S-adenosylmethionine decarboxylase involved in polyamine
	biosynthesis. It is essential for polyamine homeostasis, and normal plant
	embryogenesis, growth and development.
Synonyms:	SAMDC, S-ADENOSYLMETHIONINE DECARBOXYLASE
Immunogen:	KLH-conjugated synthetic peptide (15 aa from central section) derived from
	Arabidopsis thaliana SAMDC (AT3G02470).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
	Peptide affinity form antibody available upon request at <u>info@phytoab.com</u> .
Reconstitution:	Reconstitution with 150 µl of sterile water.
	"Note: please spin tube briefly prior to opening it to avoid any losses that might
	occur from lyophilized material adhering to the cap or sides of the tube".
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Storage:	12 months from date of receipt, -20 to -70°C as supplied.
	6 months, -20 to -70°C under sterile conditions after reconstitution.
	1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the
	temperature recommended above.

## **Application Information**


<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected/apparent MW:	40 kDa
Predicted Reactivity:	For more species homologues information, please contact tech

PhytoAB Inc.



#### support at tech@phytoab.com.

### **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.
Signals from the degraded recombinant protein were also detected.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.
Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).
Detection: using chemiluminescence substrate and image were captured with CCD camera.

Research Use Only