

## Anti-ERECTA-LIKE 2, C terminal antibody

Catalog: PHY2009A

## **Product Information**

Description:	Rabbit polyclonal antibody	
Background:	ERL2 is an ERECTA-family receptor kinase that regulates stomatal	
	patterning and inflorescence development. It suppresses stomatal formation in	
	leaves by perceiving EPF2 peptides, ensuring proper stomatal spacing. ERL2	
	also controls inflorescence architecture by promoting compact growth and floral	
	organogenesis.	
Synonyms:	ERL2, ERECTA-LIKE 2	
Immunogen:	KLH-conjugated synthetic peptide (18 aa from C terminal section) derived from	
	Arabidopsis thaliana ERL2 (AT5G07180).	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Immunogen affinity purified	
Reconstitution:	Reconstitution with 150 $\mu$ l of sterile 1XPBS (PH=7.4) containing 30% glycerol.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70°C as supplied.	
	6 months, -20 to -70°C under sterile conditions after reconstitution.	
	1 month, 2 to 8°C under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the	
	temperature recommended above.	


## **Application Information**

<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected / apparent MW:	107 kDa
Predicted Reactivity:	For more species homologues information, please contact tech
	support at <u>tech@phytoab.com</u> .

lesearch Use Only



## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.
Electrophoresis: 12% SDS-PAGE
Transfer: blotting to NC (nitrocellulose) membrane for 1 h.
Blocking: 5% skim milk at RT or 4°C for 1 h.
Primary antibody: 1:1000 dilution overnight at 4°C.

**Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.



**Research Use Only**