

## Anti-BRAHMA-INTERACTING PROTEIN 1, C-terminal antibody

Catalog: PHY7692A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** BRIP1 is a core component of SWI/SNF complexes. It interacts with BRM and

may be responsible for ensuring proper complex assembly and association with

chromatin.

Synonyms: BRIP1, BRAHMA-INTERACTING PROTEIN 1

**Immunogen:** KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Arabidopsis thaliana BRIP1 (AT3G03460).

Form: Lyophilized

**Quantity:** 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of sterile 1XPBS (PH=7.4) containing 30% glycerol.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## Application Information

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.


Expected / apparent MW: 41 kDa

Predicted Reactivity: For more species homologues information, please contact tech

support at tech@phytoab.com.



## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 46 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.