

Anti-Phototropin 2 antibody

Catalog: PHY0232A

Product Information

Description: Rabbit polyclonal antibody

Background: PHOT2 is a membrane-bound protein serine/threonine kinase that functions as

blue light photoreceptor in redundancy with PHO1. It is involved in stomatal

opening, chloroplast movement and phototropism.

Synonyms: PHOT2, ATPHOT2, NON PHOTOTROPIC HYPOCOTYL1-LIKE, NPL1

Immunogen: KLH-conjugated synthetic peptide (16 aa from central section) derived from

Arabidopsis thaliana PHOT2 (AT5G58140).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of sterile 1×PBS (PH=7.4) containing 30% glycerol.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.


Expected / apparent MW: 102 kDa

Predicted Reactivity: For more species homologues information, please contact tech

support at tech@phytoab.com.



Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 48 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.