

# Anti-OsATG8A/B/C, C-terminal antibody

Catalog: PHY4410A

### **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** ATG8A/B/C are ubiquitin-like proteins essential for autophagosome formation

and cargo recruitment. They conjugate with phosphatidylethanolamine (PE) to anchor on autophagosomal membranes, facilitating membrane expansion, substrate recognition (via LIR motifs), and selective degradation of damaged organelles/pathogens. ATG8 isoforms also regulate autophagosome-lysosome fusion and modulate stress responses (e.g., ER/mitochondrial quality control).

**Synonyms:** ATG8A/B/C, OsATG8a/b/c

**Immunogen:** KLH-conjugated synthetic peptide (16 aa from C terminal section but not the

cleavage region of ATG4) derived from Oryza sativa ATG8A (Os07g0512200),

ATG8B (Os04g0624000) and ATG8C (Os08g0191600).

Form: Lyophilized

**Quantity**: 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 μl of sterile 1×PBS (PH=7.4) containing 30% glycerol.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

# Application Information

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 14 kDa




#### **Predicted Reactivity:**

Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Zea mays, Gossypium raimondii, Solanum tuberosum, Brassica rapa, Brassica napus, Hordeum vulgare, Vitis vinifera, Glycine max, Arabidopsis thaliana, Nicotiana tabacum, Solanum lycopersicum, Populus trichocarpa, Setaria viridis, Medicago truncatula, Panicum virgatum, Spinacia oleracea, Cucumis sativus, Sorghum bicolor, Triticum aestivum.* 

For more species homologues information, please contact tech support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 46 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were

captured with CCD camera.