

# Anti-bZIP transcription factor 60 antibody

Catalog: PHY7894A

### **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** BZIP60 is a bZIP transcription factor in Arabidopsis thaliana that regulates the

unfolded protein response (UPR) during endoplasmic reticulum (ER) stress. It is primarily located on the ER membrane in its inactive form. Under ER stress, it is spliced by IRE1, releasing its active form, which translocates to the nucleus to activate UPR target genes involved in protein folding, plant immunity, and

abiotic stress tolerance (e.g., salt, drought, cold).

Synonyms: BZIP60, ATBZIP60, BASIC REGION/LEUCINE ZIPPER MOTIF 60, BBASIC

REGION/LEUCINE ZIPPER MOTIF 60

**Immunogen:** KLH-conjugated synthetic peptide (15 aa from central section) derived from

Arabidopsis thaliana BZIP60 (AT1G42990).

Form: Lyophilized

**Quantity:** 150 μg

**Purification:** Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 μl of sterile 1XPBS (PH=7.4) containing 30% glycerol.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 33 kDa


Research Use Only



#### **Predicted Reactivity:**

For more species homologues information, please contact tech support at <a href="mailto:tech@phytoab.com">tech@phytoab.com</a>.

### **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 60 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.