

Anti-NDUFS8 antibody

Catalog: PHY1101A

Product Information

Description: Rabbit polyclonal antibody

Background: Complex I is the largest protein complex of the oxidative phosphorylation system

in mitochondrial and it catalyzes NADH-quinone oxidoreduction. It represents the main entrance site for electrons into the respiratory electron transfer chain. In Arabidopsis, Complex I have at least 49 subunits and NDUFS8

(AT1G79010/AT1G16700) is one of the subunits.

Synonyms: NDUFS8

Immunogen: KLH-conjugated synthetic peptide (14 aa from Central section) derived from

Arabidopsis thaliana NDUFS8A (AT1G79010) and NDUFS8B (AT1G16700).

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 μl of sterile 1XPBS (PH=7.4) containing 30% glycerol.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 26 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in Brassica



rapa, Solanum tuberosum, Oryza sativa, Brassica napus, Setaria viridis, Glycine max, Sorghum bicolor, Medicago truncatula, Chlamydomonas reinhardtii, Solanum lycopersicum, Gossypium raimondii, Populus trichocarpa, Triticum aestivum, Hordeum vulgare, Vitis vinifera, Zea mays, Panicum virgatum, Cucumis sativus, and 80-99% homologues with the sequence in Nicotiana tabacum, Spinacia oleracea.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 47 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.