

# Anti-NF-YC2, C-terminal antibody

Catalog: PHY7780A

## Product Information


|                                 |                                                                                                                                                                                                                                                                               |
|---------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Description:</b>             | Rabbit polyclonal antibody                                                                                                                                                                                                                                                    |
| <b>Background:</b>              | NF-YC2 is a subunit of the Nuclear Factor Y complex. It regulates abscisic acid (ABA) signaling and drought stress response by interacting with NF-YA/NF-YB and binding to CCAAT-box elements in target gene promoters, affecting seed germination and stress tolerance.      |
| <b>Synonyms:</b>                | NF-YC2, ATHAP5B, HAP5B, NUCLEAR FACTOR Y, SUBUNIT C2                                                                                                                                                                                                                          |
| <b>Immunogen:</b>               | KLH-conjugated synthetic peptide (13 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> NF-YC2 (AT1G56170).                                                                                                                                                 |
| <b>Form:</b>                    | Lyophilized                                                                                                                                                                                                                                                                   |
| <b>Quantity:</b>                | 150 µg                                                                                                                                                                                                                                                                        |
| <b>Purification:</b>            | Immunogen affinity purified                                                                                                                                                                                                                                                   |
| <b>Reconstitution:</b>          | Reconstitution with 150 µl of sterile 1XPBS (PH=7.4) containing 30% glycerol.<br>"Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".                                |
| <b>Stability &amp; Storage:</b> | Use a manual defrost freezer and avoid repeated freeze-thaw cycles.<br>12 months from date of receipt, -20 to -70°C as supplied.<br>6 months, -20 to -70°C under sterile conditions after reconstitution.<br>1 month, 2 to 8°C under sterile conditions after reconstitution. |
| <b>Shipping:</b>                | The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.                                                                                                                                                                       |

## Application Information

|                                |                                                                                                                                  |
|--------------------------------|----------------------------------------------------------------------------------------------------------------------------------|
| <b>Recommended Dilution:</b>   | Western Blot (1:1000-1:2000)<br>Note: Optimal dilutions/concentrations should be determined by the end user.                     |
| <b>Expected / apparent MW:</b> | 23 kDa                                                                                                                           |
| <b>Predicted Reactivity:</b>   | For more species homologues information, please contact tech support at <a href="mailto:tech@phytoab.com">tech@phytoab.com</a> . |

Research Use Only

## Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 52 kDa.

**Electrophoresis:** 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

**Secondary antibody:** 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.