

## Anti-MYB46, C-terminal antibody

Catalog: PHY7743A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** MYB46 is a master transcriptional regulator. It directly activates the biosynthesis

of secondary cell wall components (cellulose, xylan, lignin) by controlling downstream transcription factors and enzyme genes, thereby defining

secondary wall thickness and strength in xylem vessels and fibers.

**Synonyms:** MYB46, ATMYB46, MYB DOMAIN PROTEIN 46

**Immunogen:** KLH-conjugated synthetic peptide (19 aa from C terminal section) derived from

Arabidopsis thaliana MYB46 (AT5G12870).

Form: Lyophilized

**Quantity**: 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of sterile 1XPBS (PH=7.4) containing 30% glycerol.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.


Expected / apparent MW: 32 kDa

Predicted Reactivity: For more species homologues information, please contact tech

support at tech@phytoab.com.



## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 55 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.