

# **Anti-TCP24 antibody**

Catalog: PHY8055S

### **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** TCP24 is a TCP family transcription factor. It plays a pivotal role in the control of

morphogenesis of shoot organs by negatively regulating the expression of boundary-specific genes such as CUC genes, probably through the induction of

miRNA (e.g. miR164).

Synonyms: TCP24, ATTCP24, TEOSINTE BRANCHED 1, CYCLOIDEA, AND PCF FAMILY

24

**Immunogen:** KLH-conjugated synthetic peptide (17 aa from central section) derived from

Arabidopsis thaliana TCP24 (AT1G30210).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## Application Information

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.


**Expected / apparent MW:** 36 kDa

Predicted Reactivity: For more species homologues information, please contact tech



### support at tech@phytoab.com.

### **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.