

## **Anti-Argonaute 5 antibody**

Catalog: PHY0603S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** Ago proteins are ubiquitously expressed and bind to siRNAs or miRNAs to

guide post-transcriptional gene silencing either by destabilization of the mRNA or by translational repression. AGO5 is involved in miRNA (microRNA)-directed

target cleavage.

**Synonyms:** AGO5, ATAGO5

Immunogen: KLH-conjugated synthetic peptide (17 aa from N terminal section) derived from

Arabidopsis thaliana AGO5 (AT2G27880).

Form: Lyophilized

**Quantity**: 150 μg **Purification**: Serum

Peptide affinity form antibody available upon request at <a href="mailto:info@phytoab.com">info@phytoab.com</a>.

**Reconstitution:** Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.


Expected/apparent MW: 111 kDa

**Predicted Reactivity:** For more species homologues information, please contact tech

support at tech@phytoab.com.



## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide

for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or  $4^{\circ}$ C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.