

## Anti-Plasma membrane H<sup>+</sup>-ATPase antibody

Catalog: PHY2380A

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** H<sup>+</sup>-ATPase is a primary proton pump in plants. It generates an electrochemical

gradient across the plasma membrane, which drives the secondary active transport of nutrients (e.g., K<sup>+</sup>, NO<sub>3</sub><sup>-</sup>) and regulates cell elongation, stomatal

opening, and pH homeostasis, crucial for growth and stress responses.

**Synonyms**: H<sup>+</sup>-ATPase

**Immunogen:** KLH-conjugated synthetic peptide (17 aa from C terminal section) derived from

Arabidopsis thaliana AHA4 (AT3G47950).

Form: Lyophilized

**Quantity**: 150 μg

Purification: Immunogen affinity purified

**Reconstitution:** Reconstitution with 150 µl of sterile 1XPBS (PH=7.4) containing 30% glycerol.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

**Expected / apparent MW:** 106 kDa

**Predicted Reactivity:** Among species analyzed, the sequence of the synthetic peptide used


for immunization is 80-99% homologues with the sequence in

Gossypium raimondii, Cucumis sativus, Glycine max.



For more species homologues information, please contact tech support at <a href="tech@phytoab.com">tech@phytoab.com</a>.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 48 kDa.

Electrophoresis: 12% SDS-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP)

(Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.