

Anti-Guanine nucleotide-binding protein alpha-1 subunit, C-terminal antibody

Catalog: PHY4177S

Product Information

Description:	Rabbit polyclonal antibody
Background:	GPA1 is the guanine nucleotide-binding protein alpha-1 subunit (GP-alpha-1). It is involved in morphogenesis, Gibberellin signal transduction, R gene-mediated disease resistance, and Brassinosteroid (BR) response.
Synonyms:	Os05g0333200, d1, GPA1, GA1, OsGA1, RGA1, dwf1, XA7, RGA, D1/RGA1, D89, TGW5, OsTGW5, SRG5, OsSRG5
Immunogen:	KLH-conjugated synthetic peptide (14 aa from C terminal section) derived from <i>Oryza sativa</i> Os05g0333200.
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum
Reconstitution:	Reconstitution with 150µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	44 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used

Research Use Only

for immunization is 100% homologues with the sequence in *Zea mays*, *Triticum aestivum*, *Hordeum vulgare*, *Sorghum bicolor*, *Setaria viridis*, *Panicum virgatum*, and 80-99% homologues with the sequence in *Populus trichocarpa*, *Glycine max*, *Gossypium raimondii*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.