

Anti-CYCLINC1-1 antibody

Catalog: PHY8005S

Product Information

Description: Rabbit polyclonal antibody

Background: CYCLINC1-1 is a cyclin protein that functions as a key negative regulator of

abscisic acid (ABA) signaling. It interacts with and inhibits the transcription factor

ABI5, thereby modulating ABA responses and repressing seed germination.

Synonyms: CYCLINC1-1, CYCC1-1

Immunogen: KLH-conjugated synthetic peptide (16 aa from N-terminal section) derived from

Arabidopsis thaliana CYCLINC1-1 (AT5G48640).

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

Reconstitution: Reconstitution with 150 µl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected / apparent MW: 30 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used

for immunization is 87% homologues with the sequence in Nicotiana


tabacum, Gossypium raimondii, Cucumis sativus, Vitis vinifera,



Populus trichocarpa, Triticum aestivum, and 81% homologues with the sequence in Spinacia oleracea, Hordeum vulgare, Brassica napus, Brassica rapa, Zea mays.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 46 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4° C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured

with CCD camera.