

## Anti-NADH dehydrogenase subunit 24 kD, mitochondrial antibody

Catalog: PHY1088S

## **Product Information**

**Description:** Rabbit polyclonal antibody

**Background:** Complex I is the largest protein complex of the oxidative phosphorylation system

in mitochondrial and it catalyzes NADH-quinone oxidoreduction. Complex I represents the main entrance site for electrons into the respiratory electron transfer chain. In Arabidopsis, Complex I have at least 49 subunits and 24 kD

(AT4G02580) is one of the subunit.

**Synonyms:** NADH-ubiquinone oxidoreductase 24 kDa subunit, 24 kD, 24 kDa subunit

**Immunogen:** KLH-conjugated synthetic peptide (16 aa from C terminal section) derived from

Arabidopsis thaliana AT4G02580.

Form: Lyophilized

Quantity:150 μgPurification:Serum

Peptide affinity form antibody available upon request at info@phytoab.com.

**Reconstitution:** Reconstitution with 150 μl of sterile water.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

**Stability &**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

**Storage:** 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

**Shipping:** The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

## **Application Information**

**Recommended Dilution:** Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.



Expected / apparent MW: 28 kDa

Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used


for immunization is 100% homologues with the sequence in *Brassica* napus, *Brassica rapa*, *Medicago truncatula*, *Glycine max*, and 92% homologues with the sequence in *Populus trichocarpa*, *Vitis vinifera*,

Cucumis sativus, Gossypium raimondii, Triticum aestivum.

For more species homologues information, please contact tech

support at tech@phytoab.com.

## **Application Example**



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 40 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L

(HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured

with CCD camera.