

Anti-ELONGATED HYPOCOTYL 5 antibody

Catalog: PHY1908

Product Information

Description:	Mouse monoclonal antibody
Background:	HY5 is a basic leucine zipper (bZIP) transcription factor with nuclear localization. It is involved in light-regulated transcriptional activation of G-box-containing promoters. HY5 plays a role in anthocyanin accumulation in far-red light and blue light, but not in red light or in the dark.
Synonyms:	HY5, ELONGATED HYPOCOTYL 5, REVERSAL OF THE DET PHENOTYPE 5, TED 5
Immunogen:	Recombinant protein of HY5 (1-168 aa) derived from <i>Arabidopsis thaliana</i> AT5G11260.
Form:	Lyophilized
Quantity:	150 µg
Purification:	Protein A purified
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4) containing 30% glycerol. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	18 kDa
Confirmed Reactivity:	<i>Arabidopsis thaliana</i>
Predicted Reactivity:	For more species homologues information, please contact tech

Research Use Only

support at tech@phytoab.com.

Application Example

Example1:



PHY1908

Electrophoresis: 15% SDS-Urea-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.


Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:5000 dilution using Goat Anti-Mouse IgG H&L (HRP) (Cat# PHY6006).

Detection: using chemiluminescence substrate and image were captured with CCD camera.

Example2:



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 28 kDa.

Electrophoresis: 15% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:5000 dilution using Goat Anti- Mouse IgG H&L(HRP) (Cat# PHY6006).

Detection: using chemiluminescence substrate and image were captured with CCD camera.