

Anti-TIC214 antibody

Catalog: PHY1625A

Product Information

Description:	Rabbit polyclonal antibody
Background:	TIC214/YCF1 are chloroplast-encoded membrane proteins. As a key downstream target of the RNA-binding protein PBR1, YCF1 is essential for the coordinated biogenesis of Photosystem I, cytochrome <i>b₆f</i> , and NADH dehydrogenase complexes. It is believed to act as a scaffold or platform during the assembly and stabilization of these photosynthetic complexes.
Synonyms:	TIC214, TRANSLOCON AT THE INNER ENVELOPE MEMBRANE OF CHLOROPLASTS 214, YCF1.2
Immunogen:	KLH-conjugated synthetic peptide (18 aa from C terminal section) derived from <i>Arabidopsis thaliana</i> TIC214 (ATCG01130).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4) containing 30% glycerol. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	214 kDa
Predicted Reactivity:	The sequence of the synthetic peptide used for immunization is 80-

Research Use Only

99% homologues with the sequence in *Brassica napus* and *Brassica rapa*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 43 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.