

Anti-OsSSIIIa antibody

Catalog: PHY4655S

Product Information

Description:	Rabbit polyclonal antibody
Background:	OsSSIIIa (Os08g0191433/FLO5) is a chloroplastic/amyloplastic soluble starch synthase that plays an important role in generating relatively long amylopectin chains in the rice endosperm. Its mutation leads to a white-core floury endosperm (flo5) phenotype, increased resistant starch (RS) content, and reduced amylopectin long chains.
Synonyms:	OsSSIIIa, SSIII-2, OsSSIII-2, OsSS3A, SS3A, SS3a, OsSSSIIIa, SSIIIa, Flo5
Immunogen:	KLH-conjugated synthetic peptide (17 aa from central section) derived from <i>Oryza sativa</i> OsSSIIIa (Os08g0191433).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	211 kDa

Research Use Only

Predicted Reactivity:

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 40 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.