

Anti-OsbZIP46 antibody

Catalog: PHY4405A

Product Information

Description:	Rabbit polyclonal antibody
Background:	OsbZIP46 is a nuclear bZIP transcription factor that acts as a positive regulator of ABA signaling and drought stress tolerance. It is involved in the response to water deprivation and salt stress, and its expression is induced by drought, heat, hydrogen peroxide, and ABA.
Synonyms:	BZIP46, ABL1, bZIP46, OsABL1, OsbZIP46, OsAREB8, OsABF2, OsABF3, ABF2, ABF3, OsbZIP46CA1, OsZIP46, ZIP46, OsbZIP46/OsABF2
Immunogen:	KLH-conjugated synthetic peptide (16 aa from central section) derived from <i>Oryza sativa</i> BZIP46 (Os06g0211200).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4) containing 30% glycerol. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	35 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in <i>Sorghum</i>

Research Use Only

bicolor, *Panicum virgatum*, *Zea mays*, *Setaria viridis*, and 80-99% homologues with the sequence in *Triticum aestivum*, *Hordeum vulgare*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 50 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.