

Anti-CBP20 antibody

Catalog: PHY1334A

Product Information

Description:	Rabbit polyclonal antibody
Background:	CBP20 is the small subunit of the nuclear cap-binding complex (CBC). It binds the 5' cap of pre-mRNAs and pri-miRNAs, functioning in RNA metabolism, pre-mRNA splicing, and miRNA processing. It localizes to the nucleus and the nuclear cap-binding complex.
Synonyms:	CBP20, ATCBP20, CAP-BINDING PROTEIN 20
Immunogen:	KLH-conjugated synthetic peptide (16 aa from N terminal section) derived from <i>Arabidopsis thaliana</i> CBP20 (AT5G44200)
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of sterile 1XPBS (PH=7.4) containing 30% glycerol. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information


Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	30 kDa
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in <i>Brassica napus</i> , <i>Brassica rapa</i> , and 80-99% homologues with the sequence in

Research Use Only

Oryza sativa, *Setaria viridis*, *Panicum virgatum*, *Zea mays*, *Sorghum bicolor*, *Gossypium raimondii*, *Spinacia oleracea*, *Solanum tuberosum*, *Glycine max*, *Solanum lycopersicum*, *Nicotiana tabacum*, *Triticum aestivum*, *Hordeum vulgare*, *Populus trichocarpa*, *Vitis vinifera*, *Medicago truncatula*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng, 10 ng and 25 ng recombinant protein containing the peptide for immunization and having a molecular mass of 45 kDa.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.