Anti-Histone H4 antibody

Catalog: PHY2317A

Product Information

Description:	Rabbit polyclonal antibody
Background:	H4 is histone 4.
Synonyms:	H4, HIS4, HISTONE H4
Immunogen:	KLH-conjugated synthetic peptide (16 aa from Central section) derived from *Arabidopsis thaliana* (AT2G28740, AT5G59970, AT5G59690, AT3G53730, AT3G46320, AT3G45930, AT1G07820, AT1G07660).
Form:	Lyophilized
Quantity:	150 µg
Purification:	Immunogen affinity purified
Reconstitution:	Reconstitution with 150 µl of 0.01 M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70℃ as supplied. 6 months, -20 to -70℃ under sterile conditions after reconstitution. 1 month, 2 to 8℃ under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4℃. Upon receipt, store it immediately at the temperature recommended above.


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected / apparent MW:	11 kDa
Confirmed Reactivity:	Coming soon
Predicted Reactivity:	Among species analyzed, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in *Oryza sativa*, *Panicum virgatum*, *Zea mays*, *Glycine max*, *Brassica rapa*, *Hordeum vulgare*, *Cucumis sativus*, *Medicago truncatula*, *Vitis vinifera*, *Triticum aestivum*, *Populus trichocarpa*, *Gossypium raimondii*, *Nicotiana*

tabacum, *Spinacia oleracea*, *Sorghum bicolor*, *Physcomitrium patens*, *Setaria viridis*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



Recom: 2.5 ng and 15 ng recombinant protein containing the peptide for immunization and having a molecular mass of 46 kDa.

Electrophoresis: 12% SDS-PAGE.

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4℃ for 1 h.

Primary antibody: 1:1000 dilution overnight at 4℃.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.