

Anti-Glutathione synthase 2 antibody

Catalog: PHY0135A

Product Information

Description: Rabbit polyclonal antibody

Background: GSH2 is similar to glutathione synthetases, which catalyzes one of the early

steps in glutathione biosynthesis. Two transcripts have been detected; the

longer transcript is less abundant and the protein is localized to the chloroplast.

The smaller transcript, in which the transit peptide is truncated, is localized to

the cytosol.

Synonyms: GSH2, ATGSH2, GLUTATHIONE SYNTHETASE 2, GSHB

Immunogen: KLH-conjugated synthetic peptide of GSH2 derived from Arabidopsis thaliana

AT5G27380.

Form: Lyophilized

Quantity: 150 μg

Purification: Immunogen affinity purified

Reconstitution: Reconstitution with 150 µl of 0.01 M sterile PBS.

"Note: please spin tube briefly prior to opening it to avoid any losses that might

occur from lyophilized material adhering to the cap or sides of the tube".

Stability &Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

Storage: 12 months from date of receipt, -20 to -70°C as supplied.

6 months, -20 to -70°C under sterile conditions after reconstitution.

1 month, 2 to 8°C under sterile conditions after reconstitution.

Shipping: The product is shipped at 4°C. Upon receipt, store it immediately at the

temperature recommended above.

Application Information

Recommended Dilution: Western Blot (1:1000-1:2000)

Note: Optimal dilutions/concentrations should be determined by the

end user.

Expected/apparent MW: 60 / 56 kDa

Confirmed Reactivity: Arabidopsis thaliana


Predicted Reactivity: Among species analyzed, the sequence of the synthetic peptide used



for immunization is 100% homologues with the sequence in *Populus* trichocarpa, Vitis vinifera, Gossypium raimondii, Cucumis sativus, Medicago truncatula, Glycine max, Brassica rapa, Brassica napus, and 80-99% homologues with the sequence in Oryza sativa Japonica Group, Hordeum vulgare, Nicotiana tabacum, Solanum tuberosum, Solanum lycopersicum.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



PHY0135A

10 µl total chloroplast protein from Arabidopsis thaliana leaf.

Electrophoresis: 12% SDS-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP)

(Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with

CCD camera.