

Anti-Endochitinase CHI antibody

Catalog: PHY1514S

Product Information

Description:	Rabbit polyclonal antibody	
Background:	SO2 pre-treatment significantly enhanced the activities of defence-related	
	enzymes including phenylalanine ammonia-lyase (PAL), polyphenol oxidase	
	(PPO), β -1,3-glucanase (BGL) and chitinase (CHI).	
Synonyms:	СНІ	
Immunogen:	KLH-conjugated synthetic peptide of CHI derived from Arabidopsis thaliana	
	AT2G43570.	
Form:	Lyophilized	
Quantity:	150 µg	
Purification:	Serum	
	Peptide affinity form antibody available upon request at info@phytoab.com.	
Reconstitution:	Reconstitution with 150 µl of sterile water.	
	"Note: please spin tube briefly prior to opening it to avoid any losses that might	
	occur from lyophilized material adhering to the cap or sides of the tube".	
Stability &	Use a manual defrost freezer and avoid repeated freeze-thaw cycles.	
Storage:	12 months from date of receipt, -20 to -70 $^\circ\mathrm{C}$ as supplied.	
	6 months, -20 to -70 $^\circ C$ under sterile conditions after reconstitution.	
	1 month, 2 to 8 $^\circ C$ under sterile conditions after reconstitution.	
Shipping:	The product is shipped at 4° C. Upon receipt, store it immediately at the	
	temperature recommended above.	


Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000)
	Note: Optimal dilutions/concentrations should be determined by the
	end user.
Expected/apparent MW:	30 kDa
Confirmed Reactivity:	Arabidopsis thaliana
Predicted Reactivity:	For more species homologues information, please contact tech
	support at <u>tech@phytoab.com</u> .

Research Use Only



Application Example



PHY1514S

Lane 1: 25 µg total protein from *Arabidopsis thaliana* leaf. Lane 2: 50 µg total protein from *Arabidopsis thaliana* leaf. Lane 3: 100 µg total protein from *Arabidopsis thaliana* leaf. **Electrophoresis:** 15% SDS-Urea-PAGE **Transfer:** blotting to NC (nitrocellulose) membrane for 1 h. **Blocking:** 5% skim milk at RT or 4°C for 1 h. **Primary antibody:** 1:1000 dilution overnight at 4°C.

Secondary antibody: 1:10000 dilution using Goat Anti-Rabbit IgG H&L(HRP) (Cat# PHY6000) **Detection:** using chemiluminescence substrate and image were captured with CCD camera.



Research Use Only

PhytoAB Inc.