

Anti-Sal1 phosphatase antibody

Catalog: PHY0151S

Product Information

Description:	Rabbit polyclonal antibody
Background:	Sal1 phosphatase (AT5G63980) is a bifunctional protein that has 3'(2'),5'-bisphosphate nucleotidase and inositol polyphosphate 1-phosphatase activities. It is involved in the response to cold, drought, and ABA as well as the degradation of small mRNAs. It has been shown that Sal1 phosphatase regulates light-dependent repression of hypocotyl elongation and flowering time via its 3'(2'), 5'-bisphosphate nucleotidase activity.
Synonyms:	SAL1
Immunogen:	KLH-conjugated synthetic peptide of SAL1 derived from <i>Arabidopsis thaliana</i> AT5G63980.
Form:	Lyophilized
Quantity:	150 µg
Purification:	Serum Peptide affinity form antibody available upon request at info@phytoab.com .
Reconstitution:	Reconstitution with 150 µl of sterile water. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
Shipping:	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

Application Information

Recommended Dilution:	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
Expected/apparent MW:	44 / 38 kDa

Research Use Only

Confirmed Reactivity:


Arabidopsis thaliana

Predicted Reactivity:

Among 25 analyzed species, the sequence of the synthetic peptide used for immunization is 80-99% homologues with the sequence in *Brassica napus*.

For more species homologues information, please contact tech support at tech@phytoab.com.

Application Example



PHY0151S

Lane 1: 10 µg total protein from *Arabidopsis thaliana* leaf.

Lane 2: 25 µg total protein from *Arabidopsis thaliana* leaf.

Lane 3: 50 µg total protein from *Arabidopsis thaliana* leaf.

Electrophoresis: 15% SDS-Urea-PAGE

Transfer: blotting to NC (nitrocellulose) membrane for 1 h.

Blocking: 5% skim milk at RT or 4°C for 1 h.

Primary antibody: 1:2000 dilution overnight at 4°C.

Secondary antibody: 1:20000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

Detection: using chemiluminescence substrate and image were captured with CCD camera.