

# Anti-Enolase antibody

Catalog: PHY0213

## Product Information

<b>Description:</b>	Rabbit polyclonal antibody
<b>Background:</b>	Enolases are key glycolytic enzymes that are highly conserved in prokaryotic and eukaryotic organisms and are among the most abundant cytosolic proteins.
<b>Synonyms:</b>	Enolase, ENO2, ENOLASE 2, LOS2, LOW EXPRESSION OF OSMOTICALLY RESPONSIVE GENES 2
<b>Immunogen:</b>	KLH-conjugated synthetic peptide of Enolase derived from <i>Arabidopsis thaliana</i> AT2G36530.
<b>Form:</b>	Lyophilized
<b>Quantity:</b>	150 µg
<b>Purification:</b>	Protein A purified
<b>Reconstitution:</b>	Reconstitution with 150µl of 0.01M sterile PBS. "Note: please spin tube briefly prior to opening it to avoid any losses that might occur from lyophilized material adhering to the cap or sides of the tube".
<b>Stability &amp; Storage:</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70°C as supplied. 6 months, -20 to -70°C under sterile conditions after reconstitution. 1 month, 2 to 8°C under sterile conditions after reconstitution.
<b>Shipping:</b>	The product is shipped at 4°C. Upon receipt, store it immediately at the temperature recommended above.

## Application Information


<b>Recommended Dilution:</b>	Western Blot (1:1000-1:2000) Note: Optimal dilutions/concentrations should be determined by the end user.
<b>Expected/apparent MW:</b>	48 kDa
<b>Confirmed Reactivity:</b>	<i>Arabidopsis thaliana</i>
<b>Predicted Reactivity:</b>	Among 25 analyzed species, the sequence of the synthetic peptide used for immunization is 100% homologues with the sequence in <i>Brassica rapa subsp. Oleifera</i> , <i>Solanum lycopersicum</i> , <i>Brassica napus</i> , and 80-99% homologues with the sequence in <i>Oryza sativa</i>

Research Use Only

*Japonica Group, Zea mays, Triticum aestivum, Spinacia oleracea.*

For more species homologues information, please contact tech support at [tech@phytoab.com](mailto:tech@phytoab.com).

## Application Example



**PHY0213**

Lane 1: 10 µg total protein from *Arabidopsis thaliana* leaf.

Lane 2: 25 µg total protein from *Arabidopsis thaliana* leaf.

Lane 3: 50 µg total protein from *Arabidopsis thaliana* leaf.

**Electrophoresis:** 12.5% SDS-Urea-PAGE

**Transfer:** blotting to NC (nitrocellulose) membrane for 1 h.

**Blocking:** 5% skim milk at RT or 4°C for 1 h.

**Primary antibody:** 1:1000 dilution overnight at 4°C.

**Secondary antibody:** 1:20000 dilution using Goat Anti-Rabbit IgG H&L (HRP) (Cat# PHY6000).

**Detection:** using chemiluminescence substrate and image were captured with CCD camera.